K_{PAR} : An optical property associated with ambiguous values *

Zhongping Lee

(Northern Gulf Institute, Mississippi State University, Stennis Space Center, MS 39529)

Abstract: In studies of aquatic environments, an optical property, K_{PAR} , which represents the diffuse attenuation coefficient of Photosynthetic Available Radiation (PAR), is used frequently. Because water's diffuse attenuation coefficient is highly spectral dependent and PAR is spectrally narrowing to wavelengths with less attenuation coefficients with increasing depth, this K_{PAR} is significantly depth dependent in the upper water column even for well-mixed waters. In this article, with an aim for more reliable attenuation products and more accurate description of PAR profiles, the ambiguity associated with K_{PAR} is highlighted, and the proper representation of the vertical variation of K_{PAR} is advocated.

Keywords: PAR; K_{PAR} ; ocean color remote sensing

K_{PAR} : 一个数值模糊的光学量

李忠平

(密西西比州立大学北部海湾研究所, MS 39529, USA)

摘 要: 光合有效辐射(PAR)的漫衰减系数 K_{PAR} 是水环境研究中经常使用的一个光学量。水体的漫衰减系数与波长紧密相关, 随着水体深度的不断增加, PAR 频谱收缩到具有更小衰减系数的波长处, 对上层水体或者混合均匀的水体而言, 此时的 K_{PAR} 与水深关系极为密切。为了更为准确的描述 PAR 剖面, 获取更可靠的光学衰减参数, 强调了 K_{PAR} 数值的模糊性, 倡导正确表达 K_{PAR} 的垂直变化。

关键词: 光合有效辐射; 漫衰减系数; 水色遥感

1 Introduction

In limnology and oceanography, K_{PAR} is a symbol commonly used to represent the diffuse attenuation coefficient of Photosynthetic Available Radiation (PAR). PAR is a broad band (350–700nm), spectrally integrated measurement of light intensity (often described as a flux of quanta per unit time) at a given depth (z). Knowing the intensity of PAR and its spatial variability, both horizontally and vertically, is of great importance for studies of heat transfer^[1-3] and phytoplankton dynamics in lakes^[4] and oceans^[4-7]. Since surface PAR can be well measured or modeled^[8], spatial information of K_{PAR} is then critical for the evaluation of PAR at depth for global oceans. K_{PAR} has also been used as an indicator of water quality^[9-10].

The depth dependence of PAR is commonly expressed as:

$$PAR(z) = PAR_0 e^{-K_{PAR} z} \quad (1)$$

Here PAR_0 is the PAR value just beneath the water's surface. Values of K_{PAR} may be estimated from either the surface chlorophyll concentration^[11-12] or Secchi disk depth^[13-14]. Commonly, the value of K_{PAR} in Eq.1 is presumed to be independent of depth in various studies^[15-17], and it is usually represented by the average diffuse attenuation, $\bar{K}_{PAR}$, within the euphotic zone (down to 1% of PAR_0)^[12,18]. However, it has been previously demonstrated that even for a homogeneous and well-mixed water column, the value of K_{PAR} changes significantly with depth^[12,19-20]. In this short note, along with numerical simulations, I demonstrate further the

* Received November 3, 2008; Accepted December 3, 2008. Dr. Zhongping Lee; E-mail: zplee@ngi.msstate.edu.

vertical variability of K_{PAR} by applying the various ways of calculating values of K_{PAR} . The objective is to highlight its ambiguity, and more importantly, to advocate more robust approaches/products for quantifying water quality and for modeling the subsurface PAR field.

2 Simulations and results

To illustrate the depth dependence of K_{PAR} , a common feature of apparent optical properties^[21-22], results of a numerical simulation by Hydrolight[®]^[23] are presented here. Hydrolight[®], a commercial software used by global

Figure 1 An example of vertical profiles of PAR and K_{PAR} , simulated using Hydrolight[®]

simulation. Note that $K_{PAR}(z)$ here is the diffuse attenuation coefficient at depth z , or the so-called *instantaneous* attenuation coefficient, defined as:

$$K_{PAR}(z) \equiv -\frac{d \ln(PAR(z))}{dz} \quad (2)$$

$K_{PAR}(z)$ is approximated here as:

$$K_{PAR}(z) = -\frac{\ln(PAR(z + \Delta z)) - \ln(PAR(z))}{\Delta z} \quad (3)$$

with a depth increment (Δz) of 0.1m.

For waters in this example, it is clear that $K_{PAR}(z)$ is depth dependent and it varies by three fold from the surface to the bottom of the euphotic zone (76.7m for this example). This is because that PAR measures integrated photons in the visible domain. Water and its dissolved and suspended constituents absorb and scatter photons spectrum selectively; therefore, PAR at greater depth is spectrally weighted towards bands that are least attenuated in the water column. This explains why $K_{PAR}(z)$ approaches an asymptotic minimum with increasing depth.

Because $K_{PAR}(z)$ changes with depth, an expression with greater fidelity to the physics for vertical PAR profile should be

$$PAR(z) = PAR_0 e^{-\int_0^z K_{PAR}(z) dz} \quad (4)$$

To write this expression in a simple fashion as Eq.1, the vertical variation of PAR can be expressed as

$$PAR(z) = PAR_0 e^{-\tilde{K}_{PAR}(z)z} \quad (5)$$

ocean optics community, is a powerful tool that can be used to accurately simulate 1-dimensional subsurface light field for various kinds aquatic environments. In the simulation here, the water is assumed to be well mixed with a chlorophyll concentration of 0.1mg/m^3 , a value frequently observed for world oceans^[24]. Absorption of colored dissolved organic matter (CDOM) at 440 nm is considered equivalent to that of chlorophyll^[25], and scattering of particulates is based on the model of Gordon and Morel^[26]. The sky is assumed cloudless with the sun at 30° from the zenith.

Fig.1 shows the vertical distributions of $PAR(z)$ and $K_{PAR}(z)$ calculated from the numerical

This $\tilde{K}_{\text{PAR}}(z)$, however, is no longer the instantaneous attenuation at depth z ($K_{\text{PAR}}(z)$), nor the vertically averaged value in the euphotic zone ($\bar{K}_{\text{PAR}}$). $\tilde{K}_{\text{PAR}}(z)$ is the vertical average of $K_{\text{PAR}}(z)$ between the surface and depth z :

$$\tilde{K}_{\text{PAR}}(z) \equiv -\frac{\ln(\text{PAR}(z)) - \ln(\text{PAR}_0)}{z} \quad (6)$$

Based on Eq.6, $\tilde{K}_{\text{PAR}}(z)$ can be easily derived from vertical profiles of $\text{PAR}(z)$. Because of its simplicity, the K_{PAR} values reported in the literature^[27] or data bases (e.g., Worldwide Ocean Optics Database-WOOD, SeaWiFS Bio-optical Archive and Storage System-Sea BASS) are likely calculated based on this expression. However, because instantaneous $K_{\text{PAR}}(z)$ changes with depth, $\tilde{K}_{\text{PAR}}(z)$ also differs if inconsistent ranges are used for its calculation (see Tab.1).

For the example shown in Fig.1, Table 1 presents the instantaneous PAR attenuation, $K_{\text{PAR}}(z)$, the depth-averaged PAR attenuation, $\tilde{K}_{\text{PAR}}(z)$, and the euphotic zone averaged PAR attenuation, $\bar{K}_{\text{PAR}}$. Clearly, because of the different definitions and the depth ranges used in their calculations, different K_{PAR} values (can differ by a factor of 3) could be obtained for

Tab.1 Values of various K_{PAR} of a homogeneous water body (from Hydrolight[®] simulated $\text{PAR}(z)$ profile)

Depth z (m)	$K_{\text{PAR}}(z)$ (m^{-1})	$\tilde{K}_{\text{PAR}}(z)$ (m^{-1})	$\bar{K}_{\text{PAR}}$ (m^{-1})
0.1	0.176	0.176	0.060
1.1	0.148	0.162	
5.1	0.085	0.121	
7.1	0.073	0.109	
10.1	0.064	0.097	
15.1	0.058	0.085	
20.1	0.057	0.078	
30.1	0.055	0.071	
40.1	0.054	0.067	
50.1	0.053	0.064	
60.1	0.053	0.062	
70.1	0.052	0.061	
80.1	0.052	0.060	
90.1	0.051	0.059	
100.1	0.051	0.058	

the same well-mixed water body. Note that, similarly, the diffuse attenuation coefficient of downwelling irradiance at a wavelength (K_d) also differs with ways of calculation and depths^[28-29], but its magnitude of variation is significantly smaller than that of K_{PAR} . Because of such inherent ambiguity associated with K_{PAR} , it is quite difficult to compare reported K_{PAR} values in the literature^[15-16,18] and data bases (WOOD, SeaBASS) before their definitions and depth ranges used in their calculations are explicitly provided. Furthermore, it is not surprising to see different parameterizations when depth-averaged K_{PAR} is empirically linked to either chlorophyll concentration^[11-12,30] or Secchi disk depth^[13,31], even if regional or temporal variations in bio-optical properties are assumed negligible. Consequently, for global oceans, significantly different K_{PAR} values could be generated from these different empirical relationships^[32].

Due to the large vertical variation of $K_{\text{PAR}}(z)$, vertical distribution of $\text{PAR}(z)$ by Eq.1 would be a coarse approximation if K_{PAR} is treated as a depth-independent variable. To illustrate this point, Figure 2 shows $\text{PAR}(z)$ obtained from the Hydrolight[®] simulation (used herein as a reference field) and that modeled by Eq.1 with two different depth-independent K_{PAR} values, respectively. When $\bar{K}_{\text{PAR}}$ (the averaged value within the euphotic zone, 0.060m^{-1}) is used, $\text{PAR}(z)$ value from Eq.1 matches true value for depths around the euphotic depth, but significantly overestimates PAR in the 0–30m range by as much as 40%. When $\tilde{K}_{\text{PAR}}(0-20)$ (average K_{PAR} for depth range of 0–20m, 0.078m^{-1}) is used, not only is the estimated euphotic zone depth shallower (~59.0m, a ~30% shoaling), the PAR values are overestimated in the 0–20 meter depth range, while they are

underestimated in the 40–70m range (by a factor up to 3). All these discrepancies could have significant impacts upon model simulations of heat transfer and primary production in the upper water column.

The ambiguity associated with K_{PAR} adds difficulties to the task of defining a standard remote-sensing product of K_{PAR} . First, it is not clear which K_{PAR} should be considered as the “standard” product because K_{PAR} value varies significantly with depth (also weakly sun-angle dependent); second, it is not clear if K_{PAR} values reported by different research groups follow the same definition or if they have used the same depth ranges (either geophysical depth or optical depth) for their calculations; and third, even if $\bar{K}_{\text{PAR}}$ is considered as the standard product (then requires all reported and to be reported K_{PAR} be calculated between surface and a depth where PAR is 1% of PAR_0), its value is only good to calculate PAR values around the euphotic depth (which is $4.6/\bar{K}_{\text{PAR}}$), but it does not yield accurate estimates of PAR at other depths (see Fig.2).

Figure 2 Modeled vertical profiles of $\text{PAR}(z)$ are compared with true $\text{PAR}(z)$ profile(circle with solid line)
Symbol square with dash line is for modeled $\text{PAR}(z)$ with a depth-independent K_{PAR} of 0.060 m^{-1} ; Symbol triangle with dot line is for modeled $\text{PAR}(z)$ with a depth-independent K_{PAR} of 0.078 m^{-1}

3 Conclusions

As shown in various studies^[12,33], treating K_{PAR} as a depth-independent property is not consistent with the physics of light propagation through an aquatic environment; and such K_{PAR} approach results in coarse, if not erroneous, approximation of PAR's vertical profile. In a broader perspective, these inconsistencies indicate that depth-independent K_{PAR} is not a robust candidate to be considered as a stand-alone product (in analogy to concentration of chlorophyll) for ocean color remote sensing. To accurately model or predict PAR levels in both horizontal and vertical dimensions from ocean color remote sensing, either spectrally-resolved light field^[34-35] or depth-dependent $\tilde{K}_{\text{PAR}}(z)$ (for spectrally-integrated approach) is better to be adopted^[20]. Presently $\tilde{K}_{\text{PAR}}(z)$ can be modeled from other well-defined properties or products, such as the diffuse attenuation coefficient at 490nm^[20], concentration of chlorophyll^[36-37], and the inherent optical properties^[19,38]. Separately, for the application of measuring water quality from observation of water color^[10], instead of using the ambiguous K_{PAR} , it is better to use water's inherent optical properties^[39-40] or photic depths^[41].

Acknowledgment: Financial support by NASA's Ocean Biology and Biogeochemistry is greatly appreciated. Discussions with Dr. Jason Jolliff also improved this article.

4 Reference

- [1] Kara AB, Wallcraft AJ, Hurlburt HE. Sea surface temperature sensitivity to water turbidity from simulations of the turbid Black Sea using HYCOM. *J Phys Oceanogr*, 2005, 35: 33-54.

- [2] Lewis MR, Carr M, Feldman G *et al.* Influence of penetrating solar radiation on the heat budget of the equatorial pacific ocean. *Nature*, 1990, **347**: 543-545.
- [3] Zaneveld JRV, Spinrad RW. An arc tangent model of irradiance in the sea. *J Geophys Res*, 1980, **85**: 4919-4922.
- [4] Fahnenstiel GL, Bridgeman TB, Lang GA *et al.* Phytoplankton productivity in Saginaw Bay, Lake Huron: Effects of zebra mussel (*Dreissena polymorpha*) colonization. *J Great Lakes Res*, 1995, **21**: 465-475.
- [5] Behrenfeld MJ, Falkowski PG. Photosynthetic rates derived from satellite-based chlorophyll concentration. *Limnol Oceanogr*, 1997, **42**: 1-20.
- [6] Platt T, Sathyendranath S. Oceanic primary production: estimation by remote sensing at local and regional scales. *Science*, 1988, **241**: 1613-1620.
- [7] Siegel DA, Doney SC, Yoder JA. The North Atlantic spring phytoplankton bloom and Sverdrup's critical depth hypothesis. *Science*, 2002, **296**: 730-733.
- [8] Frouin R, Lingner DW, Gautier C *et al.* A simple analytical formula to compute clear sky total and photosynthetically available solar irradiance at the ocean surface. *J Geophys Res*, 1989, **94**: 9731-9742.
- [9] Stumpf RP, Frayer ML, Durako MJ. Variations in water clarity and bottom albedo in Florida Bay from 1985 to 1997. *Estuaries*, 1999, **22**: 431-444.
- [10] Woodruff DL, Stumpf RP, Scope JA *et al.* Remote estimation of water clarity in optically complex estuarine waters. *Remote Sens Environ*, 1999, **68**: 41-52.
- [11] Mitchell BG, Holm-Hansen O. Observations and modeling of the Antarctic phytoplankton crop in relation to mixing depth. *Deep-Sea Res*, 1991, **38**: 981-1007.
- [12] Morel A. Optical modeling of the upper ocean in relation to its biogenous matter content (Case I waters). *J Geophys Res*, 1988, **93**: 10749-10768.
- [13] Buiteveld H. A model for calculation of diffuse light attenuation (PAR) and Secchi depth. *Neth J Aquat Ecol*, 1995, **29**: 55-65.
- [14] Weinberg S. Submarine daylight and ecology. *Mar Biol*, 1976, **37**: 291 - 304.
- [15] Carter CM, AH Ross, DR Schiel *et al.* In situ microcosm experiments on the influence of nitrate and light on phytoplankton community composition. *Journal of Experimental Marine Biology and Ecology*, 2005, **326**: 1-13.
- [16] RasmusE Graneli W, Wangberg SA. Optical studies in the Southern Ocean. *Deep-Sea Res(II)*, 2004, **51**: 2583-2597.
- [17] Rochford PA, Kara AB, Wallcraft AJ *et al.* Importance of solar subsurface heating in ocean general circulation models. *J Geophys Res*, 2001, **106**: 30923-30938.
- [18] Krick JTO. Light & photosynthesis in aquatic ecosystems. University Press, 1994.
- [19] Lee ZP, Du K, Arnone R *et al.* Penetration of solar radiation in the upper ocean-A numerical model for oceanic and coastal waters. *J Geophys Res*, 2005, **110**: C09019(doi:09010.01029/02004JC002780).
- [20] Zaneveld JRV, Kitchen JC, Mueller JL. Vertical structure of productivity and its vertical integration as derived from remotely sensed observations. *Limnol Oceanogr*, 1993, **38**: 1384-1393.
- [21] Kirk JTO. The vertical attenuation of irradiance as a function of the optical properties of the water. *Limnol Oceanogr*, 2003, **48**: 9-17.
- [22] Preisendorfer RW. Hydrologic optics vol. 1: introduction. National Technical Information Service. Also available on CD, Office of Naval Research, 1976.
- [23] Mobley CD. Hydrolight 3.0 Users' Guide. SRI International, 1995.
- [24] Antoine DJ, Andre M, Morel A. Oceanic primary production 2. Estimation at global scale from satellite (coastal zone color scanner) chlorophyll. *Global Biogeochemical Cycles*, 1996, **10**: 57-69.
- [25] Siegel DA, Maritorena S, Nelson NB *et al.* Global distribution and dynamics of colored dissolved and detrital organic materials. *J Geophys Res*, 2002, **107**: 3228 (doi:3210.1029/2001JC000965).
- [26] Gordon HR, Morel A. Remote assessment of ocean color for interpretation of satellite visible imagery: A review. Springer-Verlag, 1983.

- [27] Letelier RM, Karl DM, Abbott MR *et al.* Light driven seasonal patterns of chlorophyll and nitrate in the lower euphotic zone of the North Pacific Subtropical Gyre. *Limnol Oceanogr*, 2004, **49**: 508-519.
- [28] Berwald JD, Stramski CD, Mobley. Influences of absorption and scattering on vertical changes in the average cosine of the underwater light field. *Limnol Oceanogr*, 1995, **40**: 1347-1357.
- [29] Kirk Volume scattering function, average cosines, and the underwater light field. *Limnol Oceanogr*, 1991, **36**: 455-467.
- [30] Smith RC, Baker KS. The bio-optical state of ocean waters and remote sensing. *Limnol Oceanogr*, 1978, **23**: 247-259.
- [31] Montes-Hugo MA, Alvarez-Borrego S, Giles-Guzm AD. Horizontal sighting range and Secchi depth as estimators of underwater PAR attenuation in a coastal lagoon. *Estuaries*, 2003, **26**: 1302-1309.
- [32] Gattuso JP, Gentili B, Duarte CM *et al.* Light availability in the coastal ocean: impact on the distribution of benthic photosynthetic organisms and their contribution to primary production. *Biogeosciences*, 2006, **3**: 489-513.
- [33] Morel A, Gentili B. Radiation transport within oceanic (case 1) water. *J Geophys Res*, 2004, **109**: C06008, (doi:06010.01029/02003JC002259).
- [34] Liu CC, Carder KL, Miller RL *et al.* Fast and accurate model of underwater scalar irradiance. *Applied Optics*, 2002, **41**: 4962-4974.
- [35] Sathyendranath S, Platt T. The spectral irradiance field at the surface and in the interior of the ocean: a model for applications in oceanography and remote sensing. *J Geophys Res*, 1988, **93**: 9270-9280.
- [36] Morel A, Antoine D. Heating rate within the upper ocean in relation to its bio-optical state. *J of Physical Oceanography*, 1994, **24**: 1652-1665.
- [37] Ohlmann JC, Siegel D. Ocean radiant heating. Part II: Parameterizing solar radiation transmission through the upper ocean. *J of Physical Oceanography*, 2000, **30**: 1849-1865.
- [38] Barnard AH, Zaneveld JRV, Pegau WS *et al.* The determination of PAR levels from absorption coefficient profiles at 490nm. *Ciencias Marinas*, 1999, **25**: 487-507.
- [39] Gordon HR, Smith RC, Zaneveld JRV. Introduction to ocean optics Ocean Optics VI. *Proc SPIE*, 1980, **208**: 1-43.
- [40] IOCCG. Remote Sensing of Inherent Optical Properties: Fundamentals, Tests of Algorithms, and Applications. In: Lee ZP ed. Reports of the International Ocean-Colour Coordinating Group, No. 5. IOCCG, 2006: 126.
- [41] Lee ZP, Weidemann A, Kindle J *et al.* Euphotic zone depth: Its derivation and implication to ocean-color remote sensing. *J Geophys Res*, 2007, **112**: C03009(doi:03010.01029/02006JC003802).