

River water quality shaped by land–river connectivity in a changing climate

Received: 8 May 2022

Accepted: 22 December 2023

Published online: 7 March 2024

 Check for updates

Li Li¹✉, Julia L. A. Knapp^{2,9}, Anna Lintern^{3,9}, G.-H. Crystal Ng^{4,9},
Julia Perdril^{5,6,9}, Pamela L. Sullivan^{7,9} & Wei Zhi^{1,8,9}

River water quality is crucial to ecosystem health and water security, yet its deterioration under climate change is often overlooked in climate risk assessments. Here we review how climate change influences river water quality via persistent, gradual shifts and episodic, intense extreme events. Although distinct in magnitude, intensity and duration, these changes modulate the structure and hydro-biogeochemical processes on land and in rivers, hence reshaping land–river connectivity and the quality of river waters. To advance understanding of and forecasting capabilities for water quality in future climates, it is essential to perceive land and rivers as interconnected systems. It is also vital to prioritize research under climate extremes, where the dynamics of water quality often challenge existing theories and models and call for shifts in conceptual paradigms.

Humans have settled along meandering rivers and streams since the dawn of hunting and gathering societies¹. Today, over 50% of the global population lives within 3 km of a surface freshwater body, and 90% within 10 km (refs. 2,3). The survival of humans, along with the myriad forms of life sharing this Earth, hinges upon not only the rising and falling rhythm of inland flowing waters (quantity) but also the interwoven tapestry of their physical, chemical and biological composition and conditions (quality).

Given the visible nature of water levels in events such as floods and droughts, river water quantity is known to change in a warming climate. These changes have been accounted for in the global calculation of climate risks⁴. River water quality, however, is often considered as influenced more by human activities such as land use^{5,6} and less by climate change, and is therefore often overlooked in climate risk assessments. As an example, the most recent IPCC report barely discusses the risks of a changing climate on inland water quality⁴. The effects of climate change on water quality have therefore remained ‘invisible’⁷.

These ‘invisible’ impacts of climate change on river water quality directly influence the estimation of water availability, scarcity and

security^{8,9}, which quantifies and assesses risks associated with the amount of usable water, rather than simply considering the amount of available water. For example, when considering water quality variables such as water temperature and salinity, water scarcity levels can increase by a factor of 1.4 to 3 (ref. 10). Water availability is therefore inextricably linked to water quality. In fact, there is plenty of water on Earth: more than 70% of Earth’s surface is covered by ocean seawater, yet seawater cannot be used because of its high salt content.

Water quality is essential for human consumption. In industrialized modern society, drinking water originates from inland rivers, lakes and groundwater. These source waters often route through treatment facilities: sediments are filtered out; water hardness is reduced; and disease-causing microbes are removed. Water quality determines the costs of drinking water treatment¹¹. Elevated sediment and nutrient concentrations have been estimated to increase the operation and maintenance costs of water treatment facilities by 53% (ref. 11). The United States, for example, has spent about US\$5 trillion to improve surface and drinking water quality since 1972, or approximately 0.8% of its gross domestic product and an annual spending of US\$400 per

¹Department of Civil and Environmental Engineering, The Pennsylvania State University, University Park, PA, USA. ²Department of Earth Sciences, Durham University, Durham, UK. ³Department of Civil Engineering, Monash University, Clayton, Victoria, Australia. ⁴Department of Earth and Environmental Sciences, University of Minnesota-Twin Cities, Minneapolis, MN, USA. ⁵GUND Institute, University of Vermont, Burlington, VT, USA. ⁶Department of Geography and Geosciences, University of Vermont, Burlington, VT, USA. ⁷College of Earth Ocean, and Atmospheric Science, Oregon State University, Corvallis, OR, USA. ⁸The National Key Laboratory of Water Disaster Prevention, Yangtze Institute for Conservation and Development, Key Laboratory of Hydrologic-Cycle and Hydrodynamic-System of Ministry of Water Resources, Hohai University, Nanjing, China. ⁹These authors contributed equally: Julia L. A. Knapp, Anna Lintern, G.-H. Crystal Ng, Julia Perdril, Pamela L. Sullivan, Wei Zhi. ✉e-mail: lili@engr.psu.edu

Table 1 | Common water quality variables and examples of their potential ecological, health and economic effects

Common water quality variables ^a	Ecological, health and economic effects
Water temperature	High water temperature can, for example, reduce concentrations of dissolved oxygen, change pH and threaten aquatic ecosystem health.
Dissolved oxygen	Low dissolved oxygen (hypoxia) can suffocate aquatic life and lead to fish kill. Low dissolved oxygen during algal blooms has caused dead zones worldwide. Low dissolved oxygen can also mobilize chemicals such as toxic metals and phosphorus.
pH	pH is a fundamental variable that drives chemical and biological processes. Most organisms survive in only a narrow pH range. Fluctuations in pH can significantly influence water quality. For example, low pH can facilitate the mobilization of toxic metals.
Nutrients, including dissolved and particulate forms of nitrogen and phosphorus	Nutrients are essential for growth, but can limit or boost aquatic productivity, cause hypoxia and harmful algal blooms. They are often excessive in human-impacted areas such as agriculture and urban land. Elevated nitrate in drinking water can cause health problems (especially in babies). Nutrient pollution treatments have been estimated to cost >US\$2 billion annually in the United States ¹⁵³ .
Carbon: DOC, DIC and particulate organic carbon (POC)	High DOC concentrations cause brown colour in surface waters. DOC can also form carcinogenic disinfectant by-products during water treatment and mobilize toxic metals. Inorganic carbon balances pH, and often affects concentrations of cations such as calcium and magnesium (hardness). DOC, DIC and POC are sources of carbon dioxide to the atmosphere and are an important part of the global carbon cycle. DOC and POC are sometimes approximated as biochemical oxygen demand, and DIC can be approximated by alkalinity.
Toxic metals	Metals such as arsenic, mercury and lead are toxic for aquatic and human life. They often transport together with organic matter and bioaccumulate in the food chain.
Salinity, total dissolved solids, cations and anions	Salinity quantifies the total concentrations of dissolved ions, often measured by electrical conductivity or specific conductance. Salinity can originate from human activities such as irrigation and road salt application for de-icing or natural geochemical processes such as rock weathering. High-salinity waters with high total dissolved solids can precipitate as solids in soils, water pipes and facilities. Highly saline water modifies soil properties and reduces food productivity when used for irrigation. Salinity affects freshwater ecosystems and water availability for human consumption ⁸⁹ .
Turbidity	Turbidity measures the extent of light penetration in water. It reflects the concentrations of suspended particles (for example, solids and microorganisms). High turbidity reduces light availability for photosynthesis and increases water treatment costs.
Emerging contaminants	Emerging contaminants include pharmaceuticals, pesticides and industrial chemicals such as per- and polyfluoroalkyl substances ¹⁵⁴ . They can be harmful in multiple ways. For example, some are endocrine system disruptors, impacting the reproductivity of humans and aquatic life.
Microplastics	These include fragments of plastics less than 5 mm. They decay slowly and can enter organisms and biomagnify up the food chain.
Microbial contaminants	Pathogenic organisms cause disease in humans and animals. A common example is the pollution of waterways by faecal microbial contaminants such as <i>Cryptosporidium parvum</i> or <i>Salmonella</i> .

^aNote that the list and examples are not exhaustive. Concentrations (mass per volume) and loads (rates of export, in mass per time) of solutes and sediments are two important measures of water quality. Loads are calculated as the products of concentrations and river discharge (water volume per time). The Clean Water Act, the water quality regulation in the United States, establishes total maximum daily loads and maximum concentration levels in drinking water standards for many variables discussed here. The water quality inventory in the United State Geological Survey contains over 17 categories and hundreds of thousands of variables (<https://help.waterdata.usgs.gov/codes-and-parameters/parameters>), highlighting the complex nature of water quality. Based on refs. 155,156.

US citizen¹². This makes clean water arguably one of the most expensive environmental investments in the United States—more than the cost of clean air. Such access to clean water, although often taken for granted, is a privilege that should be available to everyone. Globally, about 26% of the population does not have access to safely managed drinking water services; an estimated 3 billion people are at risk of disease because of unknown water quality due to the lack of monitoring data¹³.

Water quality concerns not only domestic use but also ecosystem health, industry and agriculture. Aquatic life depends on good water quality for survival (Table 1)^{14,15}. High water temperature can reduce electricity production in thermoelectric plants¹⁶. Saline irrigation waters degrade soil properties and reduce food production¹⁷. Deteriorating water quality also impacts global carbon–climate feedback: solutes such as dissolved carbon and nutrients drive the emission of greenhouse gases from inland waters, including methane¹⁸, carbon dioxide¹⁹ and nitrous oxide²⁰.

Climate change manifests itself in two distinct forms: persistent, gradual shifts and episodic, intense extreme events. Gradual warming elevates soil temperature and green water demand (by plants) and changes total precipitation, leading to longer durations and higher frequencies of zero flow²¹. Climate change also modifies patterns of precipitation, leading to more frequent and intense events of climate extremes including floods²², droughts²³, heatwaves²⁴ and wildfires^{25,26}.

An extreme event can be defined broadly as “an episode or occurrence in which a statistically rare or unusual climatic period alters ecosystem structure and/or functions well outside the bounds of what is considered typical or normal variability”²⁷. Extreme events often change water quality rapidly and substantially, thus exacerbating water treatment demands, threatening water supply and aquatic ecosystems^{28,29}.

Existing literature has reviewed the direct impacts of gradual climate change and extreme events on river water quality^{30–35}, with a primary focus on observed changes within rivers and streams. To forecast, manage and adapt to changing water quality, it is essential to diagnose the processes and mechanisms that drive the degradation of river water quality; these processes and mechanisms often stretch beyond rivers. Here we aim to offer a mechanistic view, illustrating that river water quality is shaped not only by processes within rivers (the focus of most existing reviews) but also by their connection to the land. We highlight the changing structure and processes on the land and in rivers and the profound impacts of land–river connectivity on river water quality.

Land–river connectivity shapes river water quality
River water quality is shaped not only by the structure and processes in rivers but also by those on land. Understanding changing water quality necessitates the comprehension of the interconnected nature of




Fig. 1 | **A conceptual diagram illustrating the propagating influence of climate change on water quality via land–river connectivity.** **a**, Most precipitated water infiltrates and travels through shallow soils and deeper groundwater aquifers before ultimately emerging in rivers. The predominant flow paths differ under dry (for example, top hillslope) and wet (for example, bottom hillslope) conditions. Under dry conditions, low flow from deeper groundwater enriched in geogenic solutes such as Ca and Si (from soil and rock weathering) predominantly feeds the river, often with low water inputs leading to potentially disconnected water puddles (light and dark blue areas from shallow soil and deeper groundwater, respectively). Under wet conditions, rivers are fed more by surface runoff (sometimes carrying sediments) and shallow soil water enriched with biogenic carbon and nutrients (for example,

C, N and P, from biogeochemical transformation of organic matter). **b**, Riverine concentrations of sediments and biogenic solutes (C, N and P) therefore increase with discharge, indicating that their concentrations are typically higher under wet conditions. The opposite is true for geogenic solutes such as Ca and Si. Their riverine concentrations often increase under drier conditions when river water is dominated by deeper groundwater⁵¹. The loads of sediments and solutes generally increase with river discharge when the climate becomes wetter. Sediments and solutes are generated by soil erosion and biogeochemical reactions such as soil respiration, nutrient transformation, and chemical weathering on both land and in rivers. The rates and thermodynamics of these reactions are regulated by climate-driven temperature (T), water content (S_w) and oxygen (O_2) levels both on land and in rivers.

land and rivers. The influence of processes within rivers (in-stream processes) tends to be more important under dry and warm conditions when land inputs are minimal. As conditions become wetter, the influence of source waters from the land gains prominence^{36–38} (Fig. 1). River water quality therefore depends on the extent of land–river connectivity and the input from different flow paths at different depths shaped by distinct biogeochemical conditions^{39,40}.

Decades of water tracer analysis and global surveys has shown that the majority of the global river water derives from ‘old’ water that has routed through and interacted with the land subsurface, instead of the relatively recent ‘new’ rainfall and snowfall (within two to three months)^{41–43}. This contrasts the common perception that most river water originates directly from recent precipitation. In fact, after the arrival of precipitated water on the land surface (global river and stream surface areas account for about 0.6% of Earth’s non-glaciated land surface⁴⁴), most of it then infiltrates into soils and rocks and travels along shallow and deep flow paths, often spending years to decades before

entering rivers^{45–47} (Fig. 1). In some groundwater-fed rivers, precipitated water enters deeper paths, travelling for hundreds to thousands of years underground before re-emerging in rivers⁴⁸.

The structure of and processes on land, from the top of the trees to the bottom of groundwater aquifers, the so-called critical zone⁴⁹, therefore can have tremendous impacts on river water quality in a changing climate. In particular, as water travels through the land, it interacts with materials along its flow paths and mobilizes solids and solutes, which changes its biogeochemistry before it enters rivers (Fig. 1). The diversity and concentrations of mobilized solids and solutes depend on the chemical, physical and biological structure along its shallow and deep flow paths. Weathered and highly permeable soils typically reside in the shallow subsurface, which also harbour leaf litter and roots and living things enriched with organic matter. The decomposition of organic matter via microbe-mediated biogeochemical reactions (for example, soil respiration) generates biogenic solutes such as dissolved organic carbon (DOC) and various forms of nitrogen, which are often

enriched in the shallow subsurface and decline with depth^{50–52}. Less permeable parent rocks typically reside in the deeper subsurface below soils, delivering slower, older waters enriched with geogenic solutes such as calcium (Ca) and silica (Si) derived from chemical weathering⁵³.

Hydroclimatic conditions largely determine the depths of a river's source waters. Under dry conditions (for example, with low precipitation and high evapotranspiration; Fig. 1a, top hillslope), river waters are dominated by deeper groundwater enriched with geogenic solutes, reflecting chemical weathering of parent rocks in addition to in-stream processes^{51,53,54}. Under wet conditions (for example, with heavy rain or snowmelt; Fig. 1a, bottom hillslope), river water mostly derives from source waters that move rapidly via shallow, permeable, moist soils across large, well-connected areas of watersheds, integrating soil biogeochemical signatures along its shallow flow paths with enriched carbon and nutrient content. Such distinct source water chemistry and dominant flow paths under different hydrological regimes lead to the commonly observed concentration–discharge relationships for sediments and solutes (Fig. 1b). Specifically, sediments and biogenic carbon (C)-, nitrogen (N)- and phosphorus (P)-containing solutes often show increasing concentrations with increasing discharge (flushing), whereas geogenic solutes such as Ca and Si show the opposite dilution patterns^{51,55–57}. The loads, often quantified as products of concentrations and river discharge, typically increase with discharge regardless of concentration–discharge relationships, because discharge often rises by orders of magnitude as the system transitions from dry to wet conditions, far exceeding the typical within-an-order change for solute concentrations.

Hydrological conditions are projected to change in the future with climate. In regions and times that become wetter, climate change probably results in increased flushing (or loading) of materials and nutrients from the shallow subsurface into rivers. Where climate change increases aridity, material loads may be lower but the concentrations of water quality variables may increase due to increasing mass accumulation and intensifying in-stream processing⁵⁸.

The characteristics of the hydrological and biogeochemical structure and processes in undeveloped land can differ substantially from agricultural and urban lands. Agricultural lands have abundant legacy stores of nutrients in shallow soils and tile drainage that facilitates shallow flow^{59,60}. Urban watersheds are characterized by impervious surfaces that facilitate surface runoff, and sewer and stormwater pipes that enhance rapid subsurface flow and elevate nutrients in groundwater^{61–63}. These distinct structures in the subsurface could result in different hydrological flow paths and biogeochemical reactions, and therefore varying river water quality responses to a changing climate under diverse land-use conditions.

Impacts of persistent and gradual climate change

Land–river connectivity implies that persistent, gradual climate change influences water quality primarily in two ways. On relatively short timescales from hours to years, it directly modifies conditions and processes in rivers and on land. Over the longer timescales of decades to centuries, persistent climate change additionally alters the physical, chemical and ecological structure of rivers and land, exerting long-lasting impacts on the quality of river water.

Short-term alterations of conditions and processes

In the short term, climate change alters subsurface conditions on land and in rivers, including soil temperature (T), soil moisture (S_w) and oxygen (O_2) levels, all of which are key drivers of hydrological flow paths and reaction kinetics and thermodynamics that can change source water biogeochemistry^{64,65} (Fig. 1). For example, warming often enhances microbiological activities and rates of reactions such as soil respiration, nutrient transformation and chemical weathering⁶⁶. Variations in soil moisture regulate rates of biogeochemical reactions. In very dry soils, limited water content often slows down microbial activities

and reduces the rates of aerobic reactions that use O_2 (ref. 67). In very wet, O_2 -limited soils, microbes rely on the generally slower anaerobic reactions, including, for example, denitrification, iron reduction, sulfate reduction and methanogenesis, to obtain energy, which similarly slows down overall reactions⁶⁴. As a result, the rates of biogeochemical reactions often peak at intermediate soil moisture conditions where microbes can optimize the use of water and O_2 (ref. 65). Different types of reaction also lead to different solutes and gas products, with anaerobic reactions generally diversifying reaction products in source waters. In aquifers, lowering of groundwater tables under drier and warmer conditions promotes deeper penetration of O_2 . This enhances the oxidation of redox-sensitive bedrock such as those containing pyrite and other reduced metal-bearing minerals, which can mobilize toxic metals such as arsenic⁶⁸.

Long-term alterations of land and river structure

At decadal to centennial timescales, the physical, chemical and ecological structure of the land and rivers will evolve with climate change^{69–71}, further influencing water quality. For example, to adapt to warming, plants often modify their physiology and rooting architecture, growing roots deeper, shallower or more laterally to maximize water and nutrient acquisition^{72,73}. Soil microbes can adapt together with roots and alter the distribution and properties of organic matter^{66,74}. Alterations in root structure, soil aggregates and macropores have been documented to modify soil properties over annual to decadal timescales—much shorter than the centennial timescale that is typically expected^{75–77}. Dry river beds have been documented to function similarly to dry soil⁷⁸. Globally, warming has been attributed to the encroachment of woody shrubs into drylands and grasslands that cover nearly 40% of Earth's ice-free land surface^{79,80}. Woody shrubs modify not only water distribution and flow paths but also biogeochemistry, possibly accelerating reactions such as rock weathering, soil respiration and riverine solute export^{81,82}. Landscape units such as intermittent headwater streams and geographically isolated wetlands are essential for maintaining good water quality and can fluctuate rapidly between wet–dry and cold–hot transitions, making them particularly vulnerable to climate change³⁸. The extent of such modification depends on the types and rates of reactions, and the shifting drivers in a changing climate. Gradual climate change, therefore, can impart a persistent, complex, interacting influence on ecosystems, roots, microbes, soils and rocks, which modify flow paths and biogeochemical reactions and ultimately alter river water quality.

Deteriorating water quality during gradual climate change

While water quality changes often reflect the entangled effects of climate change and human activities, data from rivers with minimal anthropogenic activities can help differentiate the impacts of climate change. In remote Alpine lakes that integrate inputs from nearby mountain streams, decades of water chemistry data have shown concomitant increases in temperature, electrical conductivity and solute concentrations⁸³. In more than 500 US rivers with minimal human impacts, long-term mean concentrations of 16 commonly measured solutes universally increase with climate aridity (and decreasing mean river discharge)⁵⁸. Such patterns of higher mean concentrations in more arid climates are similarly observed at regional⁸⁴ to global^{85,86} scales, and have been attributed to lower water flushing capacity relative to the rates of solute production in arid climates⁵⁸. This implies that in a warming climate, as streamflow dwindles in many places, water quality may generally decline due to lower flushing capacity, even without other direct human impacts.

The changing structure and processes on land and in rivers have led to widespread deterioration of river water quality. The concentrations of DOC (Table 1) and associated 'water browning' have increased in Europe, North America and Asia, often attributed to climate warming or recovery from acid rain^{87,88}. In the United States, for example,






a Long-term mean of dissolved oxygen**b Long-term mean of water temperature**

Fig. 2 | Dissolved oxygen and water temperature across the globe. a, b, Global maps of long-term mean dissolved oxygen (DO; **a**) and water temperature (WT; **b**) in 2,975 sites from the GRQA database³⁸; each site has at least 40 data points over a minimum of five years. **c**, Mean dissolved oxygen–water temperature and DO_{sat} –water temperature correlations (2,975 sites); DO_{sat} is the solubility of O_2 (the capacity of water for dissolved O_2). **d, e**, Statistical distribution of long-term change rates in 395 data-rich sites with dissolved oxygen and water temperature data for at least 25 years. The figures generally show lower dissolved oxygen in places with higher water temperature. Rates of change for dissolved oxygen (**d**) and water temperature (**e**) were quantified using Theil–Sen slopes from the

R package *openair*; the ‘deseason’ option was used to account for potentially important seasonal influences. DO_{sat} was estimated using the ‘*calc_DO_sat*’ function from the stream Metabolizer R package¹⁵⁷. Positive and negative rates indicate increasing and decreasing trends, respectively. Only about 50% of the sites showed increasing water temperature (**e**) and decreasing dissolved oxygen (**d**), lower than the reported >70% in the United States and Central Europe rivers⁹⁷ and predominantly decreasing dissolved oxygen in the global survey³². These contrasting results highlight the complexity of climate change on water quality and data scarcity challenges in water quality.


salinity has increased substantially in 29–39% of the rivers; about 90% of US rivers have seen increasing pH since the mid-twentieth century, and the degree of these effects are influenced by climate-driven variations in runoff⁸⁹. Alkalinity, a measure of the water capacity to buffer pH changes (similar to dissolved inorganic carbon (DIC)), has increased since the 1980s in the eastern United States; and warming water may continue to exacerbate this trend⁹⁰. Water quality in the Upper Colorado basins in the United States shows strong dependence on lithology and climate (for example, precipitation)⁹¹. In glaciated regions, warming-induced glacier retreat and melting of ice sheets have accelerated the export of solutes such as phosphorus and at least doubled the rates of chemical weathering compared with rates two decades ago^{92,93}. Gradual and abrupt permafrost thaws have become hotspots for carbon and nutrient export into rivers and greenhouse gas emission^{94,95}.

The trends of changes across different regions of the world often vary, highlighting different drivers and conditions for water quality change. A global literature survey showed that water quality has

declined, improved or has no significant trends in a gradually changing climate in 56%, 31% and 13% of 956 case studies, respectively³². Water temperature and algae levels have generally increased. The concentrations of nutrients and pharmaceuticals have mostly increased, whereas other variables, including biochemical oxygen demand, salinity, suspended sediment, metals and microorganisms, have shown comparably increasing and decreasing trends³².

Dissolved oxygen showed a predominantly decreasing trend in a global survey³², possibly attributed to the major influence of water temperature⁹⁶. This is further corroborated by a recent study that used deep-learning-model-filled daily dissolved oxygen data to demonstrate widespread warming and dissolved oxygen decrease in >87% and >70% of 800 rivers in the United States and Central Europe, respectively⁹⁷. Examination of global dissolved oxygen data in the 2,975 sites from the Global River Water Quality Archive (GRQA) database⁹⁸, however, showed somewhat different proportional changes in trends. Although long-term mean dissolved oxygen concentrations are generally higher in warmer


a Fires



- Structure**
For example:
- Ecosystem destruction
 - Altered microbial community
 - Ash and debris layer
 - Soil hydrophobicity
 - Soil destabilization
- Processes**
For example:
- Decreasing transpiration
 - Increasing soil evaporation
 - Increasing discharge
 - Increasing surface runoff
 - Post-fire flushing

- Water quality**
- Water temperature ↑↑
 - Dissolved oxygen ↓↓
 - Nutrients ↑↑
 - Carbon ↑↑
 - Salinity ↑↑
 - Metals ↑↑
 - Sediments ↑↑
 - Algae bloom ↑↑


b Droughts



- Structure**
For example:
- Ecosystem destruction
 - Deepening roots
 - Altered microbial community
 - Stagnant water pools
 - Soil and riverbed cracking
 - Soil quality degradation
- Processes**
For example:
- Intensifying in-stream reactions
 - Deepening/shallowing flow paths
 - Shifting biogeochemistry
 - Post-drought flushing and erosion

- Water quality**
- Water temperature ↑↑
 - Dissolved oxygen ↓↓
 - Nutrients ↑↑
 - Carbon ↑↑
 - Salinity ↑↑
 - Metals ↑↑
 - Sediments ↑↑
 - Algae bloom ↑↑

c Storms and floods



— For example, organic matter, microbes
— For example, clay, sediments

- Structure**
For example:
- Ecosystem destruction
 - Altered microbial community
 - Landslides
 - Riverbank destruction
 - Soil erosion
- Processes**
For example:
- Shallowing of flow paths
 - Anoxic biogeochemistry
 - Flushing of legacy chemicals, sediments, sewage

- Water quality**
- Water temperature ↑↑
 - Dissolved oxygen ↓↓
 - Nutrients ↑↑
 - Carbon ↑↑
 - Salinity ↑↑
 - Metals ↑↑
 - Sediments ↑↑
 - Algae bloom ↑↑

Fig. 3 | A conceptual diagram illustrating the impacts of climate extremes on land and river structure, processes and, ultimately, water quality. a, Fires. b, Droughts. c, Storms and floods. The lightest, light and dark blue arrows represent water flow paths via ground surface, shallow soils and deeper subsurface, respectively. The upwards and downwards arrows in the right-most column indicate increasing and decreasing trends, respectively; the presence

and thickness of the arrows approximate the proportional occurrence of each trend. For fire impacts on water quality, the proportional trends are based on a meta-analysis in the United States¹¹¹ and may not have global representation. For droughts, and storms and floods, proportional trends are based on a global literature survey³².

ivers (Fig. 2a,b) and mean dissolved oxygen concentrations and solubility of oxygen correlate negatively with water temperature ($P < 0.001$; Fig. 2c)³², only about 50% of the sites showed increasing water temperature (Fig. 2e) and decreasing dissolved oxygen (Fig. 2d), lower than the reported >70% in the US and Central Europe rivers⁹⁷ and predominantly decreasing dissolved oxygen in the global survey³². It is possible that increasing water temperature may augment oxygen-producing photosynthesis, counteracting oxygen-consuming and solubility-related losses⁹⁹. Dissolved oxygen loss could also be exacerbated by land use and other watershed characteristics such that it does not directly scale with water temperature changes¹⁰⁰. For example, deoxygenation has been shown to occur most rapidly in agricultural rivers with the slowest warming rates⁹⁷. Data inconsistencies across sites probably play a role in these observed discrepancies. Some sites have only a few or no data points in some years, such that a few abnormal values can skew temporal trends. This highlights the

complex influence of climate change on water quality, calling for continued efforts to monitor rivers and to probe the role of entangled processes that drive global river water quality.

Impacts of episodic and intense climate extremes

In contrast to persistent and gradual climate change, climatic extremes such as droughts, heatwaves, wildfires, and storms and floods are episodic and intense¹⁰¹. They rapidly change land structure via processes such as landslides, soil erosion and ecosystem destruction. Such alterations can quickly modify water flow paths, the extent of land–river connectivity, and biogeochemical reactions that mobilize solutes and sediments. Different extreme events can invoke similar reactions and exert a long-lasting but distinct influence on river water quality^{29,102}. Here we focus on fires, droughts and storms as examples, but it is important to note that other extremes, such as heatwaves, have become increasingly consequential^{32,101,103}.

Fires

Wildfires have profound and lasting impacts on water quality via altering ecosystems and soil structures¹⁰⁴. Wildfire burns plants, litter and organic matter, produces ash, and changes the composition of microbial communities¹⁰⁵ (Fig. 3a). Lower post-fire water demand (transpiration) and nutrient demand also increases river discharge¹⁰⁶ and nutrient concentrations¹⁰⁷, respectively. Burning also increases soil hydrophobicity, which can reduce water infiltration into soils¹⁰⁶ and generate more overland water flow in post-fire storms. Soil hydrophobicity additionally induces post-fire soil erosion and the transport of wildfire ash into rivers, often leading to orders-of-magnitude increases in concentrations and fluxes of sediments, nutrients, carbon, major ions and metals, as documented in local^{108,109}, regional¹¹⁰ and continental¹¹¹ analyses. Concurrent with nutrient increases, altered C:N and C:P ratios have also been observed in many rivers, which can instigate enduring impacts on water quality and aquatic ecosystem health^{110,111}.

The effects of wildfire often peak immediately after a fire and gradually diminish over time, with durations varying from years to decades. In some areas, solutes such as cations and metals decrease back to their original concentrations within five years after a fire, whereas concentrations of particulate matter continue to increase¹¹⁰. Elevated sediments and nutrients in rivers have been observed to persist for more than 10 to 15 years after burning¹⁰⁴. An increasing frequency of wildfires is projected due to climate change and human activities¹¹², indicating growing and long-lasting impacts on water quality and water supply in the future²⁹.

Droughts

Droughts influence water quality directly via their impact on biogeochemical reactions in rivers^{30,32,113}. Prolonged droughts induce dry riverbeds and generate fragmented pools of stagnant water or low-flow channels with minimal water input from the surrounding land⁷⁸ (Fig. 3b). These low-flow conditions elevate temperature and prolong water residence times, which can intensify in-stream biogeochemical processing¹¹⁴. Prolonged stagnant and warm conditions also reduce air–water gas exchange and impede the replenishment of dissolved oxygen in rivers. In addition, low river flow is dominated by input from deeper groundwater with low dissolved oxygen¹¹⁵. All these conditions result in dissolved oxygen depletion, which has been observed to trigger greater occurrence of anoxic reactions that shift water quality away from non-drought conditions. Sustained stagnant water additionally can stimulate eutrophication and algal blooms, produce dangerous toxins that can sicken or kill people and animals, and create dead zones^{15,116}. A global survey indicated that concentrations of sediments, algae and most solutes (including salinity, carbon and nutrients) escalate during or after droughts^{32,116} (Fig. 3).

In addition to direct impacts within rivers, droughts have wide-ranging impacts on soil and ecosystems^{113,117}. Droughts can induce ecosystem response (for example, deepening roots and vegetation die off^{72,118}), destabilize soil and sediment structure by forming cracks^{78,113}, and modify the properties of organic matter and the composition of microbial communities⁷⁴. These changes often reduce rates of biogeochemical reactions. Flow paths often deepen during droughts as a result of low water content and deepening roots but can also become shallower post-drought due to the development of soil hydrophobicity. During post-drought storms and heavy precipitation, accumulated solutes and solids in dry riverbeds and land commonly flush out excessively, further deteriorating river water quality and exacerbating eutrophication¹¹⁶.

Storms and floods

Storms and floods influence water quality via modification of land structure and substantial export of solutes and sediments from land to rivers^{119,120}. Excessive water elevates water tables and surface water runoff, therefore promoting shallow flow paths¹²¹. Excessive surface flow can trigger landslides, collapse of riverbank and soil erosion¹²¹. These processes accelerate the mobilization of particulates and carbon- and

nitrogen-containing solutes such as DOC and nitrate from topsoil. This is particularly noticeable in agricultural areas, where large hydrological events such as storms and rain-on-snow flush out nutrients in top soils, making water quality especially vulnerable to shifting hydroclimate patterns¹²². Extreme wet conditions connect rivers to uplands that are often disconnected under non-flooding conditions, leading to disproportionately large pulses of 'stored' legacy solutes and nutrients entering rivers¹²³. In urban areas, excessive surface runoff and flow paths connected with impervious surface and urban infrastructure (for example, sewer pipes, wastewater treatment plants) flush out not only nutrients, carbon and metals but also emerging contaminants such as pharmaceuticals and microplastics.

Floods can additionally mobilize pathogens and microorganisms and spread waterborne disease^{123,124}. The occurrences of harmful algal blooms depend on the combination of nutrient levels and temperature, among other conditions, and often occur after storms (in addition to during droughts)¹²⁵. Fish kills are common after flooding, often caused by the compound, abnormal conditions, including high levels of organic materials, vegetation stress, low dissolved oxygen and fish trapping in hypoxic floodwater¹²⁶. In addition to the pulses of solutes and sediments, excessive water volumes during flooding often trigger hot moments of anoxic reactions, which can produce diverse solute and gas products that become rapidly mobilized under extreme wet conditions¹²⁷. A global survey indicated that concentrations of sediments, algae and almost all solutes increase during floods except dissolved oxygen and salinity³² (Fig. 3c).

Compound effects and extreme behaviours

Existing studies have primarily focused on the effects of individual events on water quality. Different types of extreme event, however, often occur simultaneously or consecutively with overlapping times. For example, droughts and heatwaves and wildfires often happen at similar times¹²⁸. Post-fire and post-drought floods are common^{104,109,116}. These simultaneously or consecutively occurring events can cause compound, more severe impacts on water quality than individual events alone^{129,130}. Prolonged droughts during summer heatwaves, followed by intense rainfall, have been observed to introduce large influxes of easily decomposable organic, leading to low dissolved oxygen conditions and fish kills¹²⁶. Droughts, heatwaves and fire all lead to high temperatures, low water content in soil and low dissolved oxygen in rivers—conditions that induce algae bloom and toxin release.

Frequent extreme events also bring frequent in-between transitioning conditions¹²⁷. Rapid dry–wet, hot–cold and oxic–anoxic fluctuations between extreme events often create abnormal conditions that challenge existing theories. For example, thermodynamic theories prescribing redox ladders of biogeochemical reactions have been contradicted by observations under rapidly changing conditions⁶⁴. Rates of soil respiration are expected to peak under oxic conditions. Periodic anoxic conditions, however, have been shown to sustain or even stimulate soil respiration during oxic–anoxic transitions, leading to unexpectedly high rates of soil carbon loss relative to static oxic conditions¹³¹ and possible high production rates of solutes such as DOC. Rewetting of dry soils leads to significant carbon and nutrient loss arising from sudden intensification of soil respiration^{74,132}. Methanogenesis, the process of methane production, is expected to occur under anoxic conditions. Yet it has been observed to occur faster and produce more methane in well-oxygenated dry soils than under anoxic conditions¹³³. Soil respiration rates under wet, anoxic conditions have been observed to approximate those under oxic conditions, suggesting potentially widespread underestimation of solute production under wet conditions¹³⁴.

These studies suggest that extreme events, whether single or compound, can push land and river systems beyond typical conditions for which data and knowledge exist. For example, stream solute concentrations measured at high frequency (every 5–30 minutes) suggest that solute concentrations respond to discharge variations in storm




Fig. 4 | The distinct impacts of gradual change and climate extremes on river water quality. **a**, Climate change manifests itself via gradual change (grey solid lines) and climate extremes (red dashed lines). Air temperature generally increases with time. Gradually changing precipitation leads to drier or wetter conditions, whereas climate extremes such as droughts, storms and floods cause disruption. **b**, Changing climate alters the structure of land and rivers, and the processes therein, including flow partitioning and biogeochemical reactions. It regulates temperature (T), water content (S_w) and O_2 levels, key drivers of

reaction rates and flow paths. Extreme events (dashed red arrows) may lead to rapid structure changes (for example, landslides, soil erosion, ecosystem destruction) that deviate from typical patterns under gradual changes (grey area). **c**, Concentrations and loads of representative water quality variables that generally increase (for example, C, N and P) or decrease (for example, Ca and Si) with increasing discharge. Climate extremes can lead to much larger uncertainty levels (light red shading) than those under typical range (grey shading), potentially deviating from patterns predicted by existing theories and models.

events differently compared with their responses during baseflow as identified using low-frequency measurements¹³⁵. Stream chemistry responses also vary by season, storm size and solutes. In other words, responses of water quality during events may fall outside the expectations of existing theories and models, thus challenging the ability to understand, generalize and predict water quality (Fig. 4).

Advancing understanding and forecasting capabilities

Water quality management will become increasingly challenging in a changing climate¹⁰². This is especially true with shifting extreme conditions, which have already been shown to deteriorate water quality to an extent that can threaten municipal water supplies²⁹. Forecasting water quality and near-term responses to extreme conditions will be essential for designing robust water infrastructure, making real-time management decisions and mitigating their impacts. Most existing theories and models of water quality have been developed for persistent and gradual climate change. They provide a strong basis for assessing responses to gradual climate change but will probably need further refinement for new climate regimes (Fig. 4). The effects of episodic, intense extreme events on water quality, however, have remained poorly understood such that historical dynamics cannot be relied on to predict future impacts of a changing climate^{35,136}. Data on river discharge and chemistry under extreme conditions are generally limited, which stymies capacity to develop new theories and models^{136,137}. It is therefore vital to direct future research efforts to (1) understand water quality response to climate extremes and (2) develop forecasting capabilities under extreme conditions.

Understanding water quality response to climate extremes

Given the poorly understood connections between extreme climate events and water quality, it is essential to explore how extreme events of different magnitude, duration and intensity influence the concentrations and loads of solutes and sediments in rivers, including thresholds

that trigger changing patterns in responses. It is also important to understand which processes become dominant during different extreme events, and how these relationships may differ from those under typical, non-extreme conditions. The co-occurrence of distinct types of extreme event also necessitates the characterization of their combined effects on water quality. For instance, droughts and heatwaves can co-occur and both increase water temperatures and reduce dissolved oxygen levels, but it is not clear which has more pronounced effects, whether their effects could compound nonlinearly, and whether compounds effects lead to thresholds or tipping points that trigger different types of response under distinct extents of land–river connectivity.

Answering these questions will require collecting data before, during and after extreme events. Data scarcity has been a long-standing challenge in the field of water quality¹³⁸. This is particularly the case under extreme conditions. For example, floods and fires often prevent manual sample collection and damage automated sensors. The episodic and intense nature of extreme conditions additionally narrows the temporal windows for data collection. Advances in technology for sturdy and robust automated sensors are critical for monitoring in extreme conditions¹³⁹. Understanding land–river connectivity also requires observations in the land subsurface, including physical and biogeochemical properties of soil and rocks and source water chemistries¹⁴⁰. These data are harder and more expensive to obtain but are essential for illuminating land processes that shape river water quality^{39,51,141}.

Beyond direct field observations, uncovering causal relationships between water quality and extreme events calls for the integration of data with process-based models¹⁴². It is likely that dominant processes under extreme conditions differ from those under baseline, typical conditions. This would require paradigm shifts in process conceptualization to build new models that integrate emerging understandings, and these models will need to undergo further testing with additional data collected under extreme conditions. Such scientific iteration between observations, models and hypothesis falsification is crucial

for discovering new knowledge and building better predictive models under climate extremes¹⁴³.

Different land and river systems often respond to climate extremes distinctly. To generalize theories and models, it is also crucial to move beyond individual sites and explore patterns and drivers across gradients of climate, land use and cover, geology, and other characteristics. This would require juxtaposition of data from different places and events, and drawing interpretations through complementary deductive and inductive approaches^{144,145}. It is possible that some datasets, especially those from high-frequency sensors, have already encoded process-relevant information under extreme conditions¹⁴⁶. Data-driven tools can be used to fill data gaps, detect patterns and identify influential factors, whereas process-based models can be leveraged to reveal dominant processes that regulate water quality responses to extremes.

Developing forecasting capabilities

Processed-based models explicitly simulate underlying hydrological and biogeochemical dynamics based on existing knowledge but suffer from limitations in representing process complexity and high computational requirements, especially in real time or for large-scale assessments. However, new advancements in modelling tools offer potentials for forecasting the future of water quality under climate change.

Machine learning models, in particular, deep-learning models with multiple hidden layers, have recently emerged as promising tools for river flow and water quality prediction¹⁴⁶. For example, long short-term memory (LSTM) neural network models have been shown to outperform traditional process-based hydrologic models, demonstrating versatility and accuracy in ungauged basins and flood forecasting. Although not yet widely used in water quality prediction, these approaches have shown promise in predicting water temperature and dissolved oxygen in rivers⁹⁶ and lakes¹⁴⁷ with no data. Most of these models, however, are trained using data under typical conditions instead of climate extremes.

Forecasting water quality during climate extremes requires model generalizability, that is, the capability to extrapolate beyond training data. Unlike process-based models, deep-learning models rely almost solely on information encoded in training data; they must see sufficient input-to-output patterns to extract trends and patterns. These limitations have inspired the development of theory- or process-guided deep learning (PGDL) that leverages the strengths of both types of model^{148,149}. In addition, emerging differentiable modelling integrates process-based equations and machine learning and can potentially support the exploration of process representations¹⁵⁰. PGDL has been shown to improve accuracy and reliability beyond training conditions and the physical realism of predictions with limited data¹⁵¹. In particular, when coupled with explainable artificial intelligence techniques, it also holds the potential to detect patterns and identify influential factors that drive water quality. In general, deep-learning models have been underused in water quality prediction and can be further explored for data filling, knowledge discovery and computational power to enhance forecasting capabilities. Data availability, however, will remain the bottleneck of forecasting under extreme conditions.

In summary, although water quality should be at the front and centre of climate adaptation¹⁵², it has been largely overlooked. Deteriorating river water quality threatens water availability and security not only for human consumption but also for aquatic ecosystem health, food and energy production, among others. The quality of river water has already shown widespread and substantial alterations due to climate change through drivers and processes on land and in rivers. These changes need to be accounted for in future climate risk assessments to avoid underestimates in water scarcity and inadequate designs in mitigation and adaptation initiatives^{8,9}. This Review particularly focuses on the influential role of land–river connectivity in regulating water quality amid gradually changing climate and episodic climate extremes. Existing models and theories serve as strong foundations

for gauging water quality responses to gradual climatic shifts but will nevertheless need further enhancements under unprecedented climate regimes. For climate extremes, forecasting water quality will necessitate fundamental paradigm shifts in process understanding and formulation of new theories and models that build upon innovative data collection technologies and strategies.

References

- Dong, G. et al. Spatial and temporal variety of prehistoric human settlement and its influencing factors in the upper Yellow River Valley, Qinghai Province, China. *J. Archaeol. Sci.* **40**, 2538–2546 (2013).
- Kummu, M., de Moel, H., Ward, P. J. & Varis, O. How close do we live to water? A global analysis of population distance to freshwater bodies. *PLoS ONE* **6**, e20578 (2011).
- Fang, Y. et al. Globally universal fractal pattern of human settlements in river networks. *Earths Future* **6**, 1134–1145 (2018).
- IPCC *Climate Change: The Physical Science Basis* (eds Masson-Delmotte, V. et al.) (Cambridge Univ. Press, 2021).
- Foley, J. A. et al. Global consequences of land use. *Science* **309**, 570–574 (2005).
- Allan, J. D. Landscapes and riverscapes: the influence of land use on stream ecosystems. *Annu. Rev. Ecol. Evol. Syst.* **35**, 257–284 (2004).
- Hannah, D. M. et al. Illuminating the ‘invisible water crisis’ to address global water pollution challenges. *Hydrol. Process* **36**, e14525 (2022).
- Liu, J. et al. Water scarcity assessments in the past, present, and future. *Earths Future* **5**, 545–559 (2017).
- Bakker, K. Water security: research challenges and opportunities. *Science* **337**, 914–915 (2012).
- van Vliet, M. T. H., Flörke, M. & Wada, Y. Quality matters for water scarcity. *Nat. Geosci.* **10**, 800–802 (2017).
- McDonald, R. I., Weber, K. F., Padowski, J., Boucher, T. & Shemie, D. Estimating watershed degradation over the last century and its impact on water-treatment costs for the world’s large cities. *Proc. Natl Acad. Sci. USA* **113**, 9117–9122 (2016).
- Keiser, D. A. & Shapiro, J. S. US water pollution regulation over the past half century: burning waters to crystal springs? *J. Econ. Perspect.* **33**, 51–75 (2019).
- Strong Systems and Sound Investments: Evidence on and Key Insights Into Accelerating Progress on Sanitation, Drinking-water and Hygiene: UN-water Global Analysis and Assessment of Sanitation and Drinking-water (GLAAS) 2022 Report* (World Health Organization, 2023).
- Woodward, G. et al. The effects of climatic fluctuations and extreme events on running water ecosystems. *Phil. Trans. R. Soc. Lond. B* **371**, 20150274 (2016).
- Lake, P. S. Ecological effects of perturbation by drought in flowing waters. *Freshw. Biol.* **48**, 1161–1172 (2003).
- van Vliet, M. T. H. et al. Vulnerability of US and European electricity supply to climate change. *Nat. Clim. Change* **2**, 676–681 (2012).
- Ortiz, A. C. et al. Dryland irrigation increases accumulation rates of pedogenic carbonate and releases soil abiotic CO₂. *Sci. Rep.* **12**, 464 (2022).
- Rocher-Ros, G. et al. Global methane emissions from rivers and streams. *Nature* **621**, 530–535 (2023).
- Raymond, P. A. et al. Global carbon dioxide emissions from inland waters. *Nature* **503**, 355–359 (2013).
- Yao, Y. et al. Increased global nitrous oxide emissions from streams and rivers in the Anthropocene. *Nat. Clim. Change* **10**, 138–142 (2020).
- Zipper, S. C. et al. Pervasive changes in stream intermittency across the United States. *Environ. Res. Lett.* **16**, 084033 (2021).

22. Arnell, N. W. & Gosling, S. N. The impacts of climate change on river flood risk at the global scale. *Climatic Change* **134**, 387–401 (2016).
23. McCabe, G. J. et al. A hydrologic perspective of major U.S. droughts. *Int. J. Climatol.* **43**, 1234–1250 (2023).
24. Trenberth, K. E. Changes in precipitation with climate change. *Clim. Res.* **47**, 123–138 (2011).
25. Bowman, D. M. J. S. et al. Human exposure and sensitivity to globally extreme wildfire events. *Nat. Ecol. Evol.* **1**, 0058 (2017).
26. Williams, A. P. et al. Observed impacts of anthropogenic climate change on wildfire in California. *Earths Future* **7**, 892–910 (2019).
27. Smith, M. D. An ecological perspective on extreme climatic events: a synthetic definition and framework to guide future research. *J. Ecol.* **99**, 656–663 (2011).
28. Robinne, F.-N. et al. Scientists' warning on extreme wildfire risks to water supply. *Hydrol. Process* **35**, e14086 (2021).
29. Williams, A. P. et al. Growing impact of wildfire on western US water supply. *Proc. Natl Acad. Sci. USA* **119**, e2114069119 (2022).
30. Whitehead, P. G., Wilby, R. L., Battarbee, R. W., Kernan, M. & Wade, A. J. A review of the potential impacts of climate change on surface water quality. *Hydrol. Sci. J.* **54**, 101–123 (2009).
31. Giri, S. Water quality prospective in twenty first century: status of water quality in major river basins, contemporary strategies and impediments: a review. *Environ. Pollut.* **271**, 116332 (2021).
32. van Vliet, M. T. H. et al. Global river water quality under climate change and hydroclimatic extremes. *Nat. Rev. Earth Environ.* **4**, 687–702 (2023).
33. Delpla, I., Jung, A. V., Baures, E., Clement, M. & Thomas, O. Impacts of climate change on surface water quality in relation to drinking water production. *Environ. Int.* **35**, 1225–1233 (2009).
34. Mishra, A., Alnahit, A. & Campbell, B. Impact of land uses, drought, flood, wildfire, and cascading events on water quality and microbial communities: A review and analysis. *J. Hydrol.* **596**, 125707 (2021).
35. Fabian, P. S., Kwon, H.-H., Vithanage, M. & Lee, J.-H. Modeling, challenges, and strategies for understanding impacts of climate extremes (droughts and floods) on water quality in Asia: a review. *Environ. Res.* **225**, 115617 (2023).
36. Wohl, E. et al. Connectivity as an emergent property of geomorphic systems. *Earth Surf. Process. Landf.* **44**, 4–26 (2019).
37. Wen, H. et al. Temperature controls production but hydrology regulates export of dissolved organic carbon at the catchment scale. *Hydrol. Earth Syst. Sci.* **24**, 945–966 (2020).
38. Lane, C. R. et al. Vulnerable waters are essential to watershed resilience. *Ecosystems* **26**, 1–28 (2023).
39. Li, L. et al. Toward catchment hydro-biogeochemical theories. *WIREs Water* **8**, e1495 (2021).
40. Knapp, J. L. A., Li, L. & Musolff, A. Hydrologic connectivity and source heterogeneity control concentration–discharge relationships. *Hydrol. Process.* **36**, e14683 (2022).
41. Jasechko, S., Kirchner, J. W., Welker, J. M. & McDonnell, J. J. Substantial proportion of global streamflow less than three months old. *Nat. Geosci.* **9**, 126–129 (2016).
42. McGuire, K. J. & McDonnell, J. J. Hydrological connectivity of hillslopes and streams: characteristic time scales and nonlinearities. *Water Resour. Res.* **46**, W10543 (2010).
43. McDonnell, J. J. et al. How old is streamwater? Open questions in catchment transit time conceptualization, modelling and analysis. *Hydrol. Process.* **24**, 1745–1754 (2010).
44. Allen, G. H. & Pavelsky, T. M. Global extent of rivers and streams. *Science* **361**, 585–588 (2018).
45. Tromp-van Meerveld, H. J. & McDonnell, J. J. Threshold relations in subsurface stormflow: 2. The fill and spill hypothesis. *Water Resour. Res.* **42**, W02411 (2006).
46. Hare, D. K., Helton, A. M., Johnson, Z. C., Lane, J. W. & Briggs, M. A. Continental-scale analysis of shallow and deep groundwater contributions to streams. *Nat. Commun.* **12**, 1450 (2021).
47. Kirchner, J. W., Benettin, P. & van Meerveld, I. Instructive surprises in the hydrological functioning of landscapes. *Annu Rev. Earth Planet Sci.* **51**, 277–299 (2023).
48. Sprenger, M. et al. The demographics of water: a review of water ages in the critical zone. *Rev. Geophys.* **57**, 800–834 (2019).
49. Brantley, S. L. et al. Crossing disciplines and scales to understand the critical zone. *Elements* **3**, 307–314 (2007).
50. Herndon, E. M. et al. Landscape heterogeneity drives contrasting concentration–discharge relationships in shale headwater catchments. *Hydrol. Earth Syst. Sci.* **19**, 3333–3347 (2015).
51. Stewart, B. et al. Streams as mirrors: reading subsurface water chemistry from stream chemistry. *Water Resour. Res.* **58**, e2021WR029931 (2022).
52. Musolff, A., Fleckenstein, J. H., Rao, P. S. C. & Jawitz, J. W. Emergent archetype patterns of coupled hydrologic and biogeochemical responses in catchments. *Geophys. Res. Lett.* **44**, 4143–4151 (2017).
53. Brantley, S. L. et al. Toward a conceptual model relating chemical reaction fronts to water flow paths in hills. *Geomorphology* **277**, 100–117 (2017).
54. Rose, L. A., Karwan, D. L. & Godsey, S. E. Concentration–discharge relationships describe solute and sediment mobilization, reaction, and transport at event and longer timescales. *Hydrol. Process.* **32**, 2829–2844 (2018).
55. Moatar, F., Abbott, B. W., Minaudo, C., Curie, F. & Pinay, G. Elemental properties, hydrology, and biology interact to shape concentration–discharge curves for carbon, nutrients, sediment, and major ions. *Water Resour. Res.* **53**, 1270–1287 (2017).
56. Zarnetske, J. P., Bouda, M., Abbott, B. W., Saiers, J. & Raymond, P. A. Generality of hydrologic transport limitation of watershed organic carbon flux across ecoregions of the United States. *Geophys. Res. Lett.* **45**, 11702–11711 (2018).
57. Zhi, W. & Li, L. The shallow and deep hypothesis: subsurface vertical chemical contrasts shape nitrate export patterns from different land uses. *Environ. Sci. Technol.* **54**, 11915–11928 (2020).
58. Li, L. et al. Climate controls on river chemistry. *Earths Future* **10**, e2021EF002603 (2022).
59. Basu, N. B. et al. Parsimonious modeling of hydrologic responses in engineered watersheds: structural heterogeneity versus functional homogeneity. *Water Resour. Res.* **46**, W04501 (2010).
60. Van Meter, K. J., Van Cappellen, P. & Basu, N. B. Legacy nitrogen may prevent achievement of water quality goals in the Gulf of Mexico. *Science* **360**, 427–430 (2018).
61. Oswald, C. J. et al. Integrating urban water fluxes and moving beyond impervious surface cover: a review. *J. Hydrol.* **618**, 129188 (2023).
62. Blaszcak, J. R., Delesantro, J. M., Urban, D. L., Doyle, M. W. & Bernhardt, E. S. Scoured or suffocated: urban stream ecosystems oscillate between hydrologic and dissolved oxygen extremes. *Limnol. Oceanogr.* **64**, 877–894 (2019).
63. Duncan, J. M., Welty, C., Kemper, J. T., Groffman, P. M. & Band, L. E. Dynamics of nitrate concentration–discharge patterns in an urban watershed. *Water Resour. Res.* **53**, 7349–7365 (2017).
64. Schlesinger, W. H. & Bernhardt, E. S. *Biogeochemistry: An Analysis of Global Change* 3rd edn (Academic Press, 2013).
65. Yan, Z. et al. A moisture function of soil heterotrophic respiration that incorporates microscale processes. *Nat. Commun.* **9**, 2562 (2018).

66. Jansson, J. K. & Hofmockel, K. S. Soil microbiomes and climate change. *Nat. Rev. Microbiol.* **18**, 35–46 (2020).
67. Stielstra, C. M. et al. Climatic and landscape influences on soil moisture are primary determinants of soil carbon fluxes in seasonally snow-covered forest ecosystems. *Biogeochemistry* **123**, 447–465 (2015).
68. Fendorf, S., Michael, H. A. & van Geen, A. Spatial and temporal variations of groundwater arsenic in South and Southeast Asia. *Science* **328**, 1123–1127 (2010).
69. Greaver, T. L. et al. Key ecological responses to nitrogen are altered by climate change. *Nat. Clim. Change* **6**, 836–843 (2016).
70. Stegen, J. C. et al. Variation in above-ground forest biomass across broad climatic gradients. *Glob. Ecol. Biogeogr.* **20**, 744–754 (2011).
71. Song, J. et al. A meta-analysis of 1,119 manipulative experiments on terrestrial carbon-cycling responses to global change. *Nat. Ecol. Evol.* **3**, 1309–1320 (2019).
72. Gupta, A., Rico-Medina, A. & Caño-Delgado, A. I. The physiology of plant responses to drought. *Science* **368**, 266–269 (2020).
73. McCormick, E. L. et al. Widespread woody plant use of water stored in bedrock. *Nature* **597**, 225–229 (2021).
74. Schimel, J. P. Life in dry soils: effects of drought on soil microbial communities and processes. *Annu Rev. Ecol. Evol. Syst.* **49**, 409–432 (2018).
75. Hirmas, D. R. et al. Climate-induced changes in continental-scale soil macroporosity may intensify water cycle. *Nature* **561**, 100–103 (2018).
76. Sullivan, P. L. et al. Embracing the dynamic nature of soil structure: a paradigm illuminating the role of life in critical zones of the Anthropocene. *Earth Sci. Rev.* **225**, 103873 (2022).
77. Ratajczak, Z., Nippert, J. B. & Collins, S. L. Woody encroachment decreases diversity across North American grasslands and savannas. *Ecology* **93**, 697–703 (2012).
78. Arce, M. I. et al. A conceptual framework for understanding the biogeochemistry of dry riverbeds through the lens of soil science. *Earth Sci. Rev.* **188**, 441–453 (2019).
79. Schreiner-McGraw, A. P. et al. Woody plant encroachment has a larger impact than climate change on dryland water budgets. *Sci. Rep.* **10**, 8112 (2020).
80. Stevens, N., Lehmann, C. E. R., Murphy, B. P. & Durigan, G. Savanna woody encroachment is widespread across three continents. *Glob. Change Biol.* **23**, 235–244 (2017).
81. Tune, A. K., Druhan, J. L., Lawrence, C. R. & Rempe, D. M. Deep root activity overprints weathering of petrogenic organic carbon in shale. *Earth Planet. Sci. Lett.* **607**, 118048 (2023).
82. Wen, H., Sullivan, P. L., Macpherson, G. L., Billings, S. A. & Li, L. Deepening roots can enhance carbonate weathering by amplifying CO₂-rich recharge. *Biogeosciences* **18**, 55–75 (2021).
83. Rogora, M. et al. Decadal trends in water chemistry of Alpine lakes in calcareous catchments driven by climate change. *Sci. Total Environ.* **708**, 135180 (2020).
84. Kerins, D. & Li, L. High dissolved carbon concentration in arid rocky mountain streams. *Environ. Sci. Technol.* **57**, 4656–4667 (2023).
85. Stewart, B. et al. Soil CO₂ controls short-term variation but climate regulates long-term mean of riverine inorganic carbon. *Glob. Biogeochem. Cycles* **36**, e2022GB007351 (2022).
86. Godsey, S. E., Hartmann, J. & Kirchner, J. W. Catchment chemostasis revisited: water quality responds differently to variations in weather and climate. *Hydrol. Process.* **33**, 3056–3069 (2019).
87. Monteith, D. T. et al. Dissolved organic carbon trends resulting from changes in atmospheric deposition chemistry. *Nature* **450**, 537–540 (2007).
88. de Wit, H. A., Ledesma, J. L. J. & Futter, M. N. Aquatic DOC export from subarctic Atlantic blanket bog in Norway is controlled by seasalt deposition, temperature and precipitation. *Biogeochemistry* **127**, 305–321 (2016).
89. Kaushal, S. S. et al. Freshwater salinization syndrome on a continental scale. *Proc. Natl Acad. Sci. USA* **115**, E574–E583 (2018).
90. Kaushal, S. S. et al. Increased river alkalization in the eastern U.S. *Environ. Sci. Technol.* **47**, 10302–10311 (2013).
91. Goldrich-Middaugh, G. M. et al. Regional drivers of stream chemical behavior: leveraging lithology, land use, and climate gradients across the Colorado River, Texas USA. *Water Resour. Res.* **58**, e2022WR032155 (2022).
92. Hawkings, J. et al. The Greenland Ice Sheet as a hot spot of phosphorus weathering and export in the Arctic. *Glob. Biogeochem. Cycles* **30**, 191–210 (2016).
93. Li, X. et al. Globally elevated chemical weathering rates beneath glaciers. *Nat. Commun.* **13**, 407 (2022).
94. Abbott, B. W. & Jones, J. B. Permafrost collapse alters soil carbon stocks, respiration, CH₄, and N₂O in upland tundra. *Glob. Change Biol.* **21**, 4570–4587 (2015).
95. Turetsky, M. R. et al. Carbon release through abrupt permafrost thaw. *Nat. Geosci.* **13**, 138–143 (2020).
96. Zhi, W., Ouyang, W., Shen, C. & Li, L. Temperature outweighs light and flow as the predominant driver of dissolved oxygen in US rivers. *Nat. Water* **1**, 249–260 (2023).
97. Zhi, W., Klinger, C., Liu, J. & Li, L. Widespread deoxygenation in warming rivers. *Nat. Clim. Change* **13**, 1105–1113 (2023).
98. Virro, H., Amatulli, G., Kmoch, A., Shen, L. & Uuemaa, E. GRQA: Global River Water Quality Archive. *Earth Syst. Sci. Data* **13**, 5483–5507 (2021).
99. Bernhardt, E. S. et al. The metabolic regimes of flowing waters. *Limnol. Oceanogr.* **63**, S99–S118 (2018).
100. Blaszczyk, J. R. et al. Extent, patterns, and drivers of hypoxia in the world's streams and rivers. *Limnol. Oceanogr. Lett.* **8**, 453–463 (2023).
101. Tassone, S. J. et al. Increasing heatwave frequency in streams and rivers of the United States. *Limnol. Oceanogr. Lett.* **8**, 295–304 (2023).
102. Khan, S. J. et al. Extreme weather events: should drinking water quality management systems adapt to changing risk profiles? *Water Res.* **85**, 124–136 (2015).
103. Woolway, R. I. et al. Lake heatwaves under climate change. *Nature* **589**, 402–407 (2021).
104. Smith, H. G., Sheridan, G. J., Lane, P. N. J., Nyman, P. & Haydon, S. Wildfire effects on water quality in forest catchments: a review with implications for water supply. *J. Hydrol.* **396**, 170–192 (2011).
105. Ice, G. G., Neary, D. G. & Adams, P. W. Effects of wildfire on soils and watershed processes. *J. For.* **102**, 16–20 (2004).
106. Ebel, B. A., Moody, J. A. & Martin, D. A. Post-fire temporal trends in soil-physical and -hydraulic properties and simulated runoff generation: insights from different burn severities in the 2013 Black Forest Fire, CO, USA. *Sci. Total Environ.* **802**, 149847 (2022).
107. Rhea, A. E., Covino, T. P. & Rhoades, C. C. Reduced N-limitation and increased in-stream productivity of autotrophic biofilms 5 and 15 years after severe wildfire. *J. Geophys. Res. Biogeosci.* <https://doi.org/10.1029/2020jg006095> (2021).
108. Murphy, S. F., McCleskey, R. B., Martin, D. A., Holloway, J. M. & Writer, J. H. Wildfire-driven changes in hydrology mobilize arsenic and metals from legacy mine waste. *Sci. Total Environ.* **743**, 140635 (2020).
109. Murphy, S. F., McCleskey, R. B., Martin, D. A., Writer, J. H. & Ebel, B. A. Fire, flood, and drought: extreme climate events alter flow paths and stream chemistry. *J. Geophys. Res. Biogeosci.* **123**, 2513–2526 (2018).

110. Rust, A. J., Saxe, S., McCray, J., Rhoades, C. C. & Hogue, T. S. Evaluating the factors responsible for post-fire water quality response in forests of the western USA. *Int. J. Wildland Fire* **28**, 769 (2019).
111. Hampton, T. B., Lin, S. & Nandita, B. Forest fire effects on stream water quality at continental scales: a meta-analysis. *Environ. Res. Lett.* **17**, 064003 (2022).
112. Abatzoglou, J. T. & Williams, A. P. Impact of anthropogenic climate change on wildfire across western US forests. *Proc. Natl Acad. Sci. USA* **113**, 11770–11775 (2016).
113. Vicente-Serrano, S. M., Quiring, S. M., Pena-Gallardo, M., Yuan, S. S. & Dominguez-Castro, F. A review of environmental droughts: increased risk under global warming? *Earth Sci. Rev.* **201**, 102953 (2020).
114. Song, X. et al. Drought conditions maximize the impact of high-frequency flow variations on thermal regimes and biogeochemical function in the hyporheic zone. *Water Resour. Res.* **54**, 7361–7382 (2018).
115. Brookfield, A. E. et al. Predicting algal blooms: are we overlooking groundwater? *Sci. Total Environ.* **769**, 144442 (2021).
116. Mosley, L. M. Drought impacts on the water quality of freshwater systems; review and integration. *Earth Sci. Rev.* **140**, 203–214 (2015).
117. Ravi, S., Breshears, D. D., Huxman, T. E. & D’Odorico, P. Land degradation in drylands: Interactions among hydrologic–aeolian erosion and vegetation dynamics. *Geomorphology* **116**, 236–245 (2010).
118. Xu, C. et al. Increasing impacts of extreme droughts on vegetation productivity under climate change. *Nat. Clim. Change* **9**, 948–953 (2019).
119. Raymond, P. A. & Saiers, J. E. Event controlled DOC export from forested watersheds. *Biogeochemistry* **100**, 197–209 (2010).
120. Wen, H. et al. From soils to streams: connecting terrestrial carbon transformation, chemical weathering, and solute export across hydrological regimes. *Water Resour. Res.* **8**, e2022WR032314 (2022).
121. Merz, B. et al. Causes, impacts and patterns of disastrous river floods. *Nat. Rev. Earth Environ.* **2**, 592–609 (2021).
122. Seybold, E. C. et al. Winter runoff events pose an unquantified continental-scale risk of high wintertime nutrient export. *Environ. Res. Lett.* **17**, 104044 (2022).
123. McGrane, S. J. Impacts of urbanisation on hydrological and water quality dynamics, and urban water management: a review. *Hydrol. Sci. J.* **61**, 2295–2311 (2016).
124. Levy, K., Woster, A. P., Goldstein, R. S. & Carlton, E. J. Untangling the impacts of climate change on waterborne diseases: a systematic review of relationships between diarrheal diseases and temperature, rainfall, flooding, and drought. *Environ. Sci. Technol.* **50**, 4905–4922 (2016).
125. Cann, K. F., Thomas, D. R., Salmon, R. L., Wyn-Jones, A. P. & Kay, D. Extreme water-related weather events and waterborne disease. *Epidemiol. Infect.* **141**, 671–686 (2013).
126. Kragh, T., Martinsen, K. T., Kristensen, E. & Sand-Jensen, K. From drought to flood: sudden carbon inflow causes whole-lake anoxia and massive fish kill in a large shallow lake. *Sci. Total Environ.* **739**, 140072 (2020).
127. Valett, H. M. et al. Biogeochemical and metabolic responses to the flood pulse in a semiarid floodplain. *Ecology* **86**, 220–234 (2005).
128. Libonati, R. et al. Drought–heatwave nexus in Brazil and related impacts on health and fires: a comprehensive review. *Ann. N. Y. Acad. Sci.* **1517**, 44–62 (2022).
129. Sutanto, S. J., Vitolo, C., Di Napoli, C., D’Andrea, M. & Van Lanen, H. A. J. Heatwaves, droughts, and fires: exploring compound and cascading dry hazards at the pan-European scale. *Environ. Int.* **134**, 105276 (2020).
130. Gruber, N., Boyd, P. W., Frölicher, T. L. & Vogt, M. Biogeochemical extremes and compound events in the ocean. *Nature* **600**, 395–407 (2021).
131. Huang, W. et al. High carbon losses from oxygen-limited soils challenge biogeochemical theory and model assumptions. *Glob. Change Biol.* **27**, 6166–6180 (2021).
132. Shumilova, O. et al. Simulating rewetting events in intermittent rivers and ephemeral streams: a global analysis of leached nutrients and organic matter. *Glob. Change Biol.* **25**, 1591–1611 (2019).
133. Angle, J. C. et al. Methanogenesis in oxygenated soils is a substantial fraction of wetland methane emissions. *Nat. Commun.* **8**, 1567 (2017).
134. Fairbairn, L. et al. Relationship between soil CO₂ fluxes and soil moisture: anaerobic sources explain fluxes at high water content. *Geoderma* **434**, 116493 (2023).
135. Knapp, J. L. A., von Freyberg, J., Studer, B., Kiewiet, L. & Kirchner, J. W. Concentration–discharge relationships vary among hydrological events, reflecting differences in event characteristics. *Hydrol. Earth Syst. Sci.* **24**, 2561–2576 (2020).
136. Brunner, M. I., Slater, L., Tallaksen, L. M. & Clark, M. Challenges in modeling and predicting floods and droughts: a review. *WIREs Water* **8**, e1520 (2021).
137. Guo, D. et al. A data-based predictive model for spatiotemporal variability in stream water quality. *Hydrol. Earth Syst. Sci.* **24**, 827–847 (2020).
138. McCabe, M. F. et al. The future of Earth observation in hydrology. *Hydrol. Earth Syst. Sci.* **21**, 3879–3914 (2017).
139. Zimmer, M. A. et al. Zero or not? Causes and consequences of zero-flow stream gage readings. *WIREs Water* **7**, e1436 (2020).
140. Holbrook, W. S. et al. Links between physical and chemical weathering inferred from a 65-m-deep borehole through Earth’s critical zone. *Sci. Rep.* **9**, 4495 (2019).
141. Li, L. et al. Expanding the role of reactive transport models in critical zone processes. *Earth Sci. Rev.* **165**, 280–301 (2017).
142. Li, L. Watershed reactive transport. *Rev. Mineral. Geochem.* **85**, 381–418 (2019).
143. Popper, K. R. *Conjectures and Refutations: The Growth of Scientific Knowledge* (Harper & Row, 1968).
144. Sivapalan, M., Blöschl, G., Zhang, L. & Vertessy, R. Downward approach to hydrological prediction. *Hydrol. Process.* **17**, 2101–2111 (2003).
145. Underwood, K. L. et al. Machine-learning reveals equifinality in drivers of stream DOC concentration at continental scales. *Water Resour. Res.* **59**, e2021WR030551 (2023).
146. Nearing, G. S. et al. What role does hydrological science play in the age of machine learning? *Water Resour. Res.* **57**, e2020WR028091 (2021).
147. Read, J. S. et al. Process-guided deep learning predictions of lake water temperature. *Water Resour. Res.* **55**, 9173–9190 (2019).
148. Appling, A. P., Oliver, S. K., Read, J. S., Sadler, J. M. & Zwart, J. A. in *Encyclopedia of Inland Waters* (eds Mehner, T. & Tockner, K.) 585–606 (Elsevier, 2022).
149. Karpatne, A. et al. Theory-guided data science: a new paradigm for scientific discovery from data. *IEEE Trans. Knowl. Data Eng.* **29**, 2318–2331 (2017).
150. Shen, C. et al. Differentiable modelling to unify machine learning and physical models for geosciences. *Nat. Rev. Earth Environ.* **4**, 552–567 (2023).

151. Willard, J. D. et al. Predicting water temperature dynamics of unmonitored lakes with meta-transfer learning. *Water. Resour. Res.* **57**, e2021WR029579 (2021).
152. Srivastava, S., Mehta, L. & Naess, L. O. Increased attention to water is key to adaptation. *Nat. Clim. Change* **12**, 113–114 (2022).
153. Dodds, W. K. et al. Eutrophication of U.S. freshwaters: analysis of potential economic damages. *Environ. Sci. Technol.* **43**, 12–19 (2009).
154. Domingo, J. L. & Nadal, M. Human exposure to per- and polyfluoroalkyl substances (PFAS) through drinking water: a review of the recent scientific literature. *Environ. Res.* **177**, 108648 (2019).
155. Berner, E. K. & Berner, R. A. *Global Environment: Water, Air and Geochemical Cycles* (Princeton Univ. Press, 2012).
156. Boyd, C. E. *Water Quality: An Introduction* (Springer Nature, 2019).
157. Appling, A. P. et al. The metabolic regimes of 356 rivers in the United States. *Sci. Data* **5**, 180292 (2018).

Acknowledgements

L.L. acknowledges the support of funding from US National Science Foundation (US NSF, EAR-2012669, 2012123, 2121621, 1911960) and US Department of Energy Environmental System Science programme (US DOE ESS, DE-SC0020146). P.L.S. acknowledges the support of funding from US NSF (EAR-2231723, 1911967) and US DOE ESS (DE-SC0023312). J.P. acknowledges the support of funding from National Science Foundation (EAR-2012123). G.-H.C.N. acknowledges the support of funding from US NSF (EAR-1759071) and US DOE ESS (DE-SC0020196). L.L. acknowledges students in the Li Reactive Water Group for feedback on an early version of the paper, and M. Wu for artistic suggestions on figures.

Author contributions

L.L. initiated the first draft and finalized the paper. All co-authors equally participated in generating ideas, creating content and editing various versions of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to Li Li.

Peer review information *Nature Climate Change* thanks Jeff Chanat and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

© Springer Nature Limited 2024